

Crop(s) Almond
Disease(s) Blossum blight and brown rot

Resistance management strategy for:

Group 2	Dicarboximides;
Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 7 + 3	SDHI + DMI;
Group 11	Quinone outside inhibitors (QoI);
Group 11+3	QoI + DMI; and
Group 11+7	QoI + SDHI.

		Maximum recommended number of sprays which can contain Group:			
		2	3	7 (inc. 7+3 & 7+11)	11 (inc. 11+3 & 7+11)
Total number of blossom blight/brown rot targeting sprays	1	1	1	1	1
	2	2	2	1	1
	3	2	2	1	2
	4	2	2	1	2
	5	2	2	1	2
	6	3	3	2	2
	7	3	3	2	2
	8	3	3	2	2
	9+	3	3	3	3

Guidelines

1. **Do not** apply consecutive sprays of products containing **Group 7** or **11**. Consecutive sprays include mixtures containing **Group 7** or **11**.
2. **Do not** apply more than three **Group 2** sprays in one season. Apply no more than two consecutive sprays before changing to another group.
3. Consecutive application includes from the end of one season to the start of the following season.
4. The spray program should be considered and the strategy applied on a whole-orchard basis.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.