

**Crop(s)**                      **Passionfruit**  
**Disease(s)**                **Alternata spot (*Alternaria* spp.)**

Resistance management strategy for:

<b>Group 2</b>	Dicarboximide;
<b>Group 7 + 3</b>	Succinate dehydrogenase inhibitor (SDHI) + DMI+ DMI;
	and
<b>Group 11</b>	Quinone outside inhibitors (QoI).

## Guidelines

1. Maintain a protective cover with a protectant fungicide such as mancozeb.
2. Limit the use of **Group 2** to strategic periods, i.e. before, during and after extended wet periods.
3. Always tank mix the **Group 2** fungicide with a protectant such as mancozeb.
4. **Do not** apply more than four **Group 2** sprays in a season.
5. The total number of **Group 7+3** and **Group 11** sprays should be no more than one-third of the total number of fungicide sprays per season.
6. **Do not** apply more than two consecutive applications of a **Group 11** spray.
7. If two consecutive applications of a **Group 7+3** or **Group 11** spray are applied, they must be followed by at least the same number of sprays from an alternative chemical group.

### Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through [www.apvma.gov.au](http://www.apvma.gov.au). The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.