

Crop(s) **Pulse crops**

Grey mould (Botrytis) and ascochyta Disease(s)

Resistance management strategy for:

Methyl benzimidazole carbamates; Group 1

Group 2 Dicarboximide;

Demethylation inhibitors (DMI) + Quinone outside Group 3 + 11

inhibitor (QoI);

Group 7* Succinate dehydrogenase inhibitors (SDHI);

SDHI + DMI: and Group 7 + 3

SDHI + phenylpyrroles (PP). Group 7 + 12

Guidelines

- 1. Not all listed products are registered for both diseases. Ensure the product is registered for the disease being targeted
- 2. Do not apply more than two Group 1, 2, 3+11, 7+3 or 7+12 sprays in one season (including seed treatment).
- 3. Do not apply more than two consecutive Group 1, 2 or 7 sprays, including from season to season and seed treatments. The final foliar spray of the previous season should be considered $\frac{53}{}$ when planning which fungicide group to use in seed treatments and the first foliar application. **Group 7** containing sprays should not make up more than one in every three of the spray program.
- 4. If a **Group 7** containing fungicide is used as a seed treatment, the first foliar fungicide used must not contain a **Group 7** fungicide.
- * Currently only available under temporary permit. Always check permits are valid prior to use.

^{*} Currently only available under temporary permit. Always check that permits are valid prior to use.