



Crop(s) Turf Disease(s) Various

Resistance management strategy for:

Group 1 Benzimidazoles; **Group 2** Dicarboxamides;

Group 3 Demethylation inhibitors (DMI);

Group 4 Phenylamides (PA);

Group 7 (N3) Succinate Dehydrogenase inhibitors (SDHI); **Group 7 + 11** SDHI + Quinone outside inhibitors (QoI);

Group 11 QoI;

Group 11 + 2 QoI + dicarboxamides;

Group 11 + 3 QoI + DMI;

Group 12 Phenylpyrroles (PP);

Group 14 Aromatic hydrocarbons (chlorophenyls and

nitroanilines);

Group 21 Quinone inside inhibitors (QiI);

Group 28 Carbamates; and Group 33 Phosphonates.

Guidelines

- Fungicide groups that are classified as medium to high risk for fungicide resistance development; Groups 1, 2, 4, 7 (N3), 11 and 21 should be rotated as a key feature to reduce the development of resistance.
- 2. **Do not** apply consecutive sprays of fungicides from the same activity group, unless mixed with a registered fungicide from a different mode of action group with no known resistance.
- 3. **Group N3** nematicides are also classified as **Group 7** fungicides and consecutive application should be avoided, even if targeting different pests/pathogens.
- 4. If consecutive sprays are applied of fungicides from a high-risk fungicide group (i.e. **Group 1**, **2**, **4**, **7**, **11** or **21**) they must be followed by at least the same number of applications of fungicide(s) from a different group(s) before the same high-risk fungicide is used again.