

Crop(s) **Wheat**
Disease(s) **Septoria blotch**

Resistance management strategy for:

Group 3	Demethylation inhibitors (DMI);
Group 7	Succinate dehydrogenase inhibitors (SDHI);
Group 3 + 7	DMI + SDHI; and
Group 11 + 3	Quinone outside inhibitors (QoI) + DMI.
Group 21	Quinone inside inhibitor (QiI).

Guidelines

1. Fungicides should be used as protectant treatments – prior to symptoms of disease when conditions are conducive to disease development.
2. In high-risk disease environments, integrated management approaches should be used to reduce fungicide resistance risk, which may include:
 - a. Removal or burning of stubble;
 - b. Crop rotation (**avoid** wheat on wheat);
 - c. Control of green bridge volunteers; and
 - d. Use of tolerant and resistant varieties.
3. **Do not** apply more than one application of a **Group 7** seed treatment with foliar activity in any two consecutive growing seasons.
4. **Do not** apply more than two applications per growing season of **Group 11** or **7** containing products. Foliar sprays and in-furrow or seed treatments, that have activity on foliar diseases, count toward total. Combinations of in-furrow and seed treatment are counted as one application.
5. If a **Group 7** seed treatment has been used with foliar activity (as determined by label claims), the first foliar fungicide applied must not contain a **Group 7** fungicide. An alternative mode of action fungicide should be applied at typical foliar fungicide timing.
6. If a **Group 7** or **11** fungicide is being applied as a foliar spray, it must be in a co-formulation or in mixture with a registered mixing partner with a different mode of action, with no known resistance. The mixture partner should also provide effective disease control when used alone on the target disease.
7. **Do not** apply more than three applications containing **Group 3** fungicides per growing season. This total of three applications includes DMIs applied as **Group 11+3** co-formulations and in-furrow or seed treatments that have activity on foliar diseases. Combinations of in-furrow and seed treatments are counted as one application.

Continued on next page.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.

8. Minimise use of **Group 3** fungicides which are known to have compromised resistance status.
9. Reduced rates, including with multiple applications, must not be used. Always used labelled rates.
10. Apply **Group 21** preventatively, using a maximum of one **Group 21** spray per crop. Do not apply **Group 21** solo, they must be applied in combination with an effective tank mix partner from a different group.

Please note:

This strategy is a guide only and does not endorse particular products, groups of products or cultural methods in terms of their performance. Always follow the product label for specific use instructions. While all effort has been made with the information supplied in this document, no responsibility, actual or implied, is taken for the day to day accuracy of product or active constituent specific information. Readers should check with the Australian Pesticides and Veterinary Medicines Authority's product database for contemporary information on products and actives. The database can be sourced through www.apvma.gov.au. The information given in this strategy is provided in good faith and without any liability for loss or damage suffered as a result of its application and use. Advice given in this strategy is valid as at 26 June 2025. All previous versions of this strategy are now invalid.

